Hémorragie du Post-Partum (HPP)

Pr Frédéric MERCIER

Département d'Anesthésie-Réanimation
Hôpital Antoine Bécclère, CLAMART
& Université de K. Bicêtre, Paris XI
(frederic.mercier@abc.aphp.fr)

Conflits d'intérêt potentiels

- ◆ Laboratoire LFB (Clottafact[®])
- ◆ Laboratoire Novonordisk (Novoseven[®])
- ◆ Laboratoire Fresenius (Voluven[®])

HPP : définition et incidence

- ◆ HPP « immédiat » : pertes sanguines > 500 ml, dans les 24 h après l'accouchement
 - ➔ ≈ 5% des accouchements
- ◆ Sévère : > 1000 ml (voire 1500 ml)
ou ↘ Hte > 10% (voire ↘ Hb > 4 g/dL)
 - ➔ ≈ 1% des accouchements

Diagnostic d'HPP : souvent pas si évident !

- ◆ **Saignement vaginal** souvent sous estimé :

- sac de recueil non utilisé

- circonscrit dans l'utérus : AU

- circonscrit derrière le placenta : HRP

- hémopéritoine : RU

- hématomes para vaginal ou pelvien : TG

- ◆ **Htc / Hb** (appareil portable) :

- varie selon le remplissage vasculaire et la vitesse du saignement

- ◆ **Instabilité hémodynamique** : trop tardive, voire absente en cas de réanimation précoce et énergique

Diagnostic d'HPP :

- ◆ Le saignement vaginal important extériorisé, la chute de l'Hb et l'hypoTA peuvent manquer
 - peut induire en erreur l'**obstétricien** !
- ◆ Critères meilleurs = signes **indirects**
 - ◆ Besoins importants de remplissage vasculaire
 - besoins de catécholamines
 - besoins de transfusion
 - ◆ Troubles de l'hémostase
 - évalués par l'**anesthésiste** !
 - **communication** cruciale avec obstétricien !

L'analyse des dossiers par le Comité National d'Experts sur la Mortalité Maternelle (CNEMM) montre que l'HPP :

- reste la 1^{ère} cause de Mortalité Maternelle (MM)
- reste la cause de MM ayant le plus fort % d'évitabilité

(1^{er}, 2nd et 3^{ème} Rapports du CNEMM : 2000, 2006, 2010)

http://www.invs.sante.fr/publications/2010/mortalite_maternelle/rapport_mortalite_maternelle.pdf

Gynécologie Obstétrique & Fertilité 33 (2005) 268–274

CONSENSUS ET RPC

Hémorragies du post-partum : recommandations du CNGOF pour la pratique clinique (décembre 2004)

Postpartum haemorrhage: recommendations for clinical practice by the CNGOF (December 2004)

F. Goffinet ^{a,b,*}, F. Mercier ^{c,d}, V. Teyssier ^{d,c}, F. Pierre ^b, M. Dreyfus ^e,
A. Mignon ^{c,d}, B. Carbonne ^b,
G. Lévy ^b, et l'ensemble du groupe de travail des RPC sur l'HPP ^l

multidisciplinaire: CNGOF, CARO-SFAR, SFR, EFS, CNSF

Journées du CNGOF, déc 2004

J Gynecol Obstet Biol Reprod 2004, vol 33, supp N°8

- 1 - Fréquence, conséquences en termes de **santé** et FDR avant l'accouchement
- 2 - Gestion du risque **anténatal**
- 3 - FDR au cours du **travail** et prévention clinique et pharmacologique de l'HPP
- 4 & 5 - Prise en charge **obstétricale / anesthésique initiale** en cas d'HPP
- 6 & 7 - Prise en charge **obstétricale / anesthésique** en cas d'une HPP qui persiste malgré les mesures initiales ou qui est **grave** d'emblée
- 8 - Modalités d'un **transfert** maternel dans le cadre d'une HPP
- 9 - Prise en charge des HPP graves : indications et techniques de **l'embolisation** artérielle
- 10 - HPP immédiat : techniques et indications de la **chirurgie**
- 11 - Gestion des **PSL** dans le cadre des HPP immédiat

Les 3 étiologies les plus fréquentes d'HPP

➔ constituent 90 à 95% de l'ensemble des HPP

Hémorragie en salle de naissance

Noter heure

Installer monitoring

Appeler aide

DA + RU

Examen sous valves
→ sutures

Vidange vésicale

Oxytocine

10-20 UI perf ± IVL
Massage utérin

Oxygène

Remplissage:
Cristalloïdes, colloïdes
± éphédrine
± taux d'Hb in situ

Vérifier 2 déter
+ RAI

2^{ème} voie ± NFS-Coag
± Vérifier
disponibilité sang

T30

Sonde urinaire
Diurèse horaire

Antibiothérapie

Réchauffement
patiente

Sulprostone:

500µg/50ml PSE
1^{ère} amp en 1h mini
± 1^{ème} amp en 5h

Bilan biologique
complet

CG pour Hb > 7g
PFC fct clinique et TP
Plaquettes si Pl < 50000
± Fibrinogène

Maintien
PAM 60-80 mm Hg

Si besoin:
noradrénaline: 0,5mg/h
à adapter

Si VVC: fémorale G

T60

**Ligatures
vasculaires**

Embolisation

Hystérectomie

(minutes)

Hémorragie en salle de naissance

Noter heure

Installer monitoring

Appeler aide

DA + RU
Examen sous valves
→ sutures

Vidange vésicale
Oxytocine
10-20 UI perf ± IVL
Massage utérin

Oxygène
Remplissage:
Cristalloïdes, colloïdes
± éphédrine
± taux d'Hb in situ

Vérifier 2 déter
+ RAI
2^{ème} voie ± NFS-Coag
± Vérifier
disponibilité sang

T30

Sonde urinaire
Diurèse horaire

Antibiothérapie

Réchauffement
patiente

Sulprostone:
500µg/50ml PSE
1^{ère} amp en 1h mini
± 1^{ème} amp en 5h

Bilan biologique
complet

CG pour Hb ≈ 9-10 g/dL
PFC:CG = 1:1
Plaquettes si Pl < 50-100000
+ Fibri si < 1,5-2g

Maintien
PAM 60-80 mm Hg

Si besoin:
noradrénaline: 0,5mg/h
à adapter

Voie centrale/écho ?
Voie artérielle ?

T60

acide
tranéxamique
(1g IV puis 3g/3h)

**Ligatures
vasculaires**

Embolisation

rFVIIa

Hystérectomie

(minutes)

Conf. Actu SFAR, sept. 2008

ELSEVIER
SAUNDERS

Anesthesiology Clin
26 (2008) 53–66

ANESTHESIOLOGY
CLINICS

Major Obstetric Hemorrhage

Frederic J. Mercier, MD, PhD^{a,*},
Marc Van de Velde, MD, PhD^b

^a*Department of Anesthesia and Intensive Care, Hopital Antoine Beclere—APHP
and Universite Paris-Sud, 157 rue de la Porte de Trivaux,
92141 Clamart Cedex BP 405, France*

^b*Department of Anaesthesiology, University Hospitals Gasthuisberg,
Katholieke Universiteit Leuven, Herestraat 49, B-3000 Leuven, Belgium*

Acide tranéxamique

(Exacyl[®])

Anti-fibrinolytic agents in post partum haemorrhage: a systematic review

Pili Ferrer, Ian Roberts*, Emma Sydenham, Karen Blackhall and Haleema Shakur

BMC Pregnancy and Childbirth 2009, **9**:29

Results: We identified three randomised controlled trials involving 461 participants. The trials compared tranexamic acid with no treatment and reported blood loss after delivery. In all three trials, allocation concealment was either inadequate or unclear. The administration of tranexamic acid was associated with a reduction in blood loss of 92 millilitres (95%CI 76 to 109). The most frequently reported adverse effect of tranexamic acid was nausea, although the increase was easily compatible with the play of chance (RR 4.63, 95%CI 0.23 to 95.14).

Study or Subgroup	Tranexamic acid			No treatment			Weight	Mean Difference IV, Fixed, 95% CI	Mean Difference IV, Fixed, 95% CI
	Mean	SD	Total	Mean	SD	Total			
Gai 2004	359.29	152.02	91	439.36	191.48	89	10.3%	-80.07 [-130.65, -29.49]	
Gohel 2007	374.92	51.46	50	472.79	43.54	50	75.8%	-97.87 [-116.55, -79.19]	
Yang 2001	243.3	106.3	94	314.8	180.9	87	13.9%	-71.50 [-115.17, -27.83]	
Total (95% CI)			235			226	100.0%	-92.37 [-108.64, -76.11]	

Heterogeneity: Chi² = 1.44, df = 2 (P = 0.49); I² = 0%
Test for overall effect: Z = 11.13 (P < 0.00001)

-23%

-200 -100 0 100 200
Favours experimental Favours control

HPP > 4-500 ml : RR = 0.44 (95%CI 0.31 to 0.64)

Essai clinique randomisé contrôlé évaluant
l'impact d'un traitement antifibrinolytique sur
le volume de l'hémorragie du post-partum

étude prospective multicentrique
EXADELI

Crit Care Med 2011 : 15(2):R117. Epub 2011 Apr 15.

Anne-Sophie DUCLOY-BOUTHORS
Maternité Jeanne de Flandre
CHRU de LILLE

→ **4g** en 1h suivi de **1g/h** pendant 6h

Efficacité

	Gr. Contrôle (n=74)	Gr. Exacyl (n=78)	<i>P</i>
↘ Hb > 4g	34	15	< 0,001
Total CG	62	28	< 0,001
Nalador [®]	36	34	NS
Embolisation	7	4	NS
Lig. / Hyst.	1 / 1	0 / 0	NS

Etude prospective multicentrique EXADELI

- ◆ réduit le volume de l'HPP de 25 à 50%
 - ◆ Mais effets secondaires à type de troubles visuels et de nausées vomissements
 - ◆ délai d'action de 30 minutes possible
- réduire les doses de moitié et administrer dès le début des troubles de l'hémostase ?

Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial

(Lancet 2010)

	Tranexamic acid (n=10 060)	Placebo (n=10 067)
Any cause of death	1463 (14.5%)	1613 (16.0%)
Bleeding	489 (4.9%)	574 (5.7%)

→ **1g** IVL suivi de **1g** en perfusion IV sur 8h

[Full Text View](#)
[Tabular View](#)
[No Study Results Posted](#)
[Related Studies](#)

World Maternal Antifibrinolytic Trial (WOMAN)

This study is currently recruiting participants.

Verified on June 2011 by London School of Hygiene and Tropical Medicine

First Received on March 30, 2009. Last Updated on June 29, 2011 [History of Changes](#)

Sponsor:	London School of Hygiene and Tropical Medicine
Collaborator:	Indian Council of Medical Research
Information provided by:	London School of Hygiene and Tropical Medicine
ClinicalTrials.gov Identifier:	NCT00872469

► Purpose

This trial is a large pragmatic randomised double-blind, placebo controlled trial to quantify the effects of the early administration of tranexamic acid on death, hysterectomy and other relevant outcomes. 15,000 adult women, after delivery who have clinically diagnosed postpartum haemorrhage, are eligible if the responsible doctor is for any reason substantially uncertain whether or not to use an antifibrinolytic agent.

Condition	Intervention	Phase
Postpartum Haemorrhage	Drug: Tranexamic acid Drug: Placebo [Saline]	Phase III

Study Type: Interventional
 Study Design: Allocation: Randomized
 Endpoint Classification: Safety/Efficacy Study
 Intervention Model: Parallel Assignment
 Masking: Double Blind (Subject, Caregiver, Investigator, Outcomes Assessor)
 Primary Purpose: Treatment

Official Title: Tranexamic Acid for the Treatment of Postpartum Haemorrhage: An International Randomised, Double Blind, Placebo Controlled Trial

Acide tranéxamique (Exacyl[®])

« Protocole Béclère » :

→ 1g IVL en 5-10 min

± 0,5 g / h (1-2 g / 2-4 h)

Fibrinogène (Clottafact[®])

RPC déc. 2004 : “L’intérêt d’un apport supplémentaire de fibrinogène dans le traitement de la CIVD est controversé”

The decrease of fibrinogen is an early predictor of the severity of postpartum haemorrhage

Charbit et al, J Thromb Haemost 2007; 5: 266-73

- ◆ La fibrinogénémie est le seul paramètre indépendant qui soit associé avec une évolution sévère de l'HPP
- ◆ Taux FI > 4 g/l = VPN 79%
- ◆ Taux FI < 2 g/l = VPP 100%
- ◆ La valeur prédictive de ces résultats s'accroissent de H0 à H4

ORIGINAL ARTICLE

Some hemostasis variables at the end of the population distributions are risk factors for severe postpartum hemorrhages

C. CHAULEUR, *† E. COCHERY-NOUVELLON, *‡ E. MERCIER, *‡§ G. AYA, ‡¶ P. FABBRO-PERAY, ** P. MISMETTI, † G. LISSADE-LAVIGNE*‡ and J.-C. GRIS*‡§

*Haematology laboratory, University Hospital, Nîmes; †EA3065, Saint-Etienne; ‡Research group EA2992, Nîmes; §Haematology laboratory, Faculty of Pharmaceutical and Biological Sciences, Montpellier 1 University, Montpellier; ¶Department of Gynaecology and Obstetrics, University Hospital, Nîmes; and **Department of biostatistics and of clinical research, University Hospital, Nîmes, France

factors. Among the 32 463 women enrolled in the NOHA First cohort, 317 developed severe PPH (S-PPH group), 1269 non-severe PPH (NS-PPH group) and the remaining individuals were considered as control women (C group). *Methods:* We performed a case-control study, including 317 triplets of women allocated from the three groups that shared the same clinical characteristics as the S-PPH group. *Results:* From values obtained 6–9 months after delivery, low (but not-deficient) levels of fibrinogen, von Willebrand factor (VWF) antigen, factor (F) XI, platelet CD42b, TRAP-induced increase of platelet CD41a and high values of serum residual prothrombin activity or closure aperture times using the collagen-ADP cartridge on the PFA-100[®] system, and blood group O, were independently associated with a significant risk of severe PPH. Being positive for at least two of these eight variables was found in 1.6%, 3.5% and 20.8% of the women from the C, the NS-PPH and the S-PPH groups, respectively, the odds ratio for S-PPH in such a case being 16.4, 95%CI (6.5–41), $P < 0.0001$.

Seuil fibri = 2 g / L

The *In Vitro* Effects of Fibrinogen Concentrate, Factor XIII and Fresh Frozen Plasma on Impaired Clot Formation After 60% Dilution

Thorsten Haas, MD*

Dietmar Fries, MD†

Corinna Velik-Salchner, MD*

Christian Reif‡

Anton Klingler, MD, PhD§

Petra Innerhofer, MD*

BACKGROUND: Previous investigations have shown that increasing fibrinogen concentration improves dilution-dependent impairment of clot formation. We conducted an *in vitro* study to explore whether substitution with fibrin-stabilizing factor XIII (FXIII) combined with fibrinogen promotes further improvement of clot formation, and whether fibrinogen administration as concentrate or fresh frozen plasma (FFP) results in comparable effects.

METHODS: Blood from six healthy donors was diluted by 60% using lactated Ringer's solution. Aliquots of diluted blood samples were incubated with two different doses of fibrinogen concentrate, FXIII concentrate, the combination of both, or with two different doses of FFP. Using thrombelastometry (ROTEM®) blood samples were analyzed at baseline (undiluted), after dilution and after supplementation. Variables were analyzed for changes from baseline, and effects of fibrinogen concentrate alone or combined with FXIII were compared with effects observed with corresponding FFP doses.

RESULTS: After 60% *in vitro* dilution of blood all ROTEM parameters and global coagulation tests changed significantly. Among the substitutes tested FXIII alone had no effect, the combination with fibrinogen improved coagulation time, α angle and fibrinogen/fibrin polymerization significantly more than did small-dose fibrinogen alone. After substituting fibrinogen, median values of all ROTEM variables were within the normal range, thereby showing dose dependency but also significant differences ($P = 0.027$) from corresponding FFP doses (EXTEM MCF FFP small dose [38 (35, 40.3) mm]), which enabled only coagulation time to be shortened to baseline levels.

CONCLUSIONS: Supplementation of fibrinogen restored all ROTEM parameters after dilution. This effect was partially enhanced by adding FXIII and was significantly stronger than for FFP substitution.

(Anesth Analg 2008;106:1360-5)

Beside assessment of fibrinogen level in PPH by thromboelastometry

Huissoud et al, BJOG 2009

CA₁₅ @ 6mm : valeur seuil pour détecter fibri < 1,5 g/l (Se = 100%, Spe = 88%)

Management of post-partum haemorrhage

J. AHONEN¹, V. STEFANOVIC² and R. LASSILA³

Departments of ¹Anaesthesia and Intensive Care, ²Obstetrics and Gynaecology and ³Internal Medicine, Division of Haematology and Laboratory Services, Coagulation Disorders, Helsinki University Hospital, Helsinki, Finland

Acta Anaesthesiol Scand 2010

risk of severe haemorrhage (Fig. 2B). This is in accordance with experimental thromboelastography findings showing that at fibrinogen 0.5 g/l, clot is not formed. Clot formation begins at 0.75 g/l, while all parameters of clot formation augment markedly from 0.75 to 3 g/l. Thus, fibrinogen is critical not only for clot strength but also to accelerate clot initiation and propagation.³⁶ However,

The use of fibrinogen concentrate to correct hypofibrinogenaemia rapidly during obstetric haemorrhage

S.F. Bell, R. Rayment,* P.W. Collins* R.E. Collis

*Department of Anaesthesia and *Department of Haematology, University Hospital of Wales, Cardiff, UK*

International Journal of Obstetric Anesthesia (2010) 19, 218–234

few reports of its use in association with continuing obstetric haemorrhage. Six cases of obstetric haemorrhage, associated with hypofibrinogenaemia, treated with fibrinogen concentrate in conjunction with platelets, fresh frozen plasma, packed red blood cells, uterotonics and obstetric intervention are described. In all cases, laboratory assessed coagulation was rapidly normalised and severe haemorrhage improved. These cases suggest that fibrinogen concentrate may be an effective addition to conventional treatments for obstetric haemorrhage associated with hypofibrinogenaemia.

Table 2 Comparison of cost and quantity of FFP, fibrinogen concentrate and cryoprecipitate required to raise plasma fibrinogen concentration by 1 g/L in a 70-kg adult

Blood product	Predicted quantity required to increase plasma fibrinogen concentration by 1 g/L (volume, mL)	Cost to increase plasma fibrinogen concentration by 1 g/L
<u>FFP</u> ⁷	4 units (<u>1000 mL</u>)	£384
Cryoprecipitate ⁷	13 units (260 mL)	£478
<u>Fibrinogen</u> concentrate ⁹	2 g (<u>100 mL</u>)	£440

Facteur VII activé recombinant (FVIIa, Novoseven[®])

Tissue factor (TF)/FVIIa, or TF/rFVIIa interaction, is necessary to initiate haemostasis

At pharmacological concentrations rFVIIa directly activates FX on the surface of locally activated platelets.

This activation will initiate the "thrombin burst" independently of FVIII and FIX. This step is independent of TF.

The thrombin burst leads to the formation of a stable clot

The use of rFVIIa in obstetric and GYN haemorrhage

Franchini et al., BJOG 2007; 114: 8-15

Table 1. The use of rFVIIa in obstetric PPH

Authors	Year	Women	Age (years)	Cause of PPH	Type of delivery	Hysterectomy	Initial dose of rFVIIa (microgram/kg)	Number of doses	Response, n (%)*
Haya <i>et al.</i> ³⁵	1998	1	28	1 FVII-Pr	1 CS	0	28	4	1
Muleo <i>et al.</i> ³⁶	1998	1	22	1 FVII-Pr	1 CS	0	20	9	1
Jimenez-Yuste <i>et al.</i> ³⁷	2000	1	30	1 FVII-Pr	1 CS	1	13.3	—**	1
Moscardo <i>et al.</i> ³⁸	2001	1	33	1 At	1 CS	1	90	9	1
Brueckner-Sabine <i>et al.</i> ³⁹	2001	1	31	1 At	1 CS	1	NR	NR	1***
Monte and Lyons ⁴⁰	2001	1	31	1 GT-Pr	1 CS	0	90	1	1
Breborowicz <i>et al.</i> ⁴¹	2002	8	33.9 (26–44)	4 At, 3 DIC, 1 Pr	7 CS, 1 VD	5/8 (62.5)	24.5 (16.7–48)	1	7/8 (87.5)
Zupancic <i>et al.</i> ⁴²	2002	1	31	1 DIC-HELLP	1 CS	0	90	1	1
Sobieszczyk <i>et al.</i> ⁴³	2002	1	29	1 IOP	1 CS	1	NR	NR	1
Eskandari <i>et al.</i> ⁴⁴	2002	1	NR	1 FVII-Pr	1 VD	0	50	2	1
Sokolic <i>et al.</i> ⁴⁵	2002	1	31	1 DIC-HELLP	1 CS	0	90	1	1
Bouwmeester <i>et al.</i> ⁴⁶	2003	1	30	1 At-Lac	1 VD	1	60	2	1
Kretzschmar <i>et al.</i> ⁴⁷	2003	1	35	1 DIC	1 CS	1	60	1	1***
Boyer-Neumann <i>et al.</i> ⁴⁸	2003	1	29	1 VWD	1 CS	0	41	5	1
Dart <i>et al.</i> ⁴⁹	2004	1	24	1 HELLP	1 VD	0	90	1	1
Boehlen <i>et al.</i> ⁵⁰	2004	1	31	1 At	1 VD	1	120	19	1
Segal <i>et al.</i> ⁵¹	2004	10	NR	3 PA; 2 At; 4 Lac; 1 UM	NR	7/10 (70.0)	88.0 (60–100)	1.1 (1–2)	10/10 (100)
Merchant <i>et al.</i> ⁵²	2004	3	30	3 HELLP-SLH	3 CS	0/3	90	2	3/3****
Kale <i>et al.</i> ⁵³	2004	1	31	1 GT-Pr	1 VD	0	36	1	1
Ahonen and Jokela ⁵⁴	2005	12	27.7 (24–37)	3 PA, 2 At, 1 PP, 6 Lac	5 CS, 7 VD	5/12 (41.7)	85.1 (42–120)	1	10/12 (83.3)
Shamsi <i>et al.</i> ⁵⁵	2005	3	30.7 (27–35)	1 PP, 2 Lac	3 CS	2/3 (66.7)	86.7 (80–90)	1	3/3 (100)
Tanchev <i>et al.</i> ⁵⁶	2005	4	NR	4 At	4 VD	0/4	72.0 (61–82)	1	4/4 (100)
Holub <i>et al.</i> ⁵⁷	2005	1	28	1 At	1 CS	1	NR	1	1
Hollnberger <i>et al.</i> ⁵⁸	2005	3	29.7 (28–31)	1 At, 1 PP, 1 At-Lac	1 CS, 2 VD	0/3	100.0 (60–120)	2	3/3
Nowacka <i>et al.</i> ⁵⁹	2005	1	30	1 IOP	1 CS	1	37.5	2	1
Verre <i>et al.</i> ⁶⁰	2006	1	24	1 At	1 CS	1	90	1	1
Palomino <i>et al.</i> ⁶¹	2006	3	NR	1 At, 1 PP, 1 placental abruption	2 CS, 1 VD	1/3 (33.3)	40	1	3/3 (100)***
Total		65	30.5 (22–44)		35 CS, 20 VD	30/65 (46.1)	65.9 (13.3–120)	1.8 (1–19)	62/65 (95.4)

**PLACE CHRONOLOGIQUE DE L'ADMINISTRATION
DU FACTEUR VII ACTIVE RECOMBINANT EN SAUVETAGE MATERNEL DANS
L'HEMORRAGIE SEVERE DU POST-PARTUM :**

Avant ou après les gestes chirurgicaux de survie :

Embolisation, ligature vasculaire et hystérectomie d'hémostase

Inclusions terminées, analyse en cours !...

Investigateur Principal :
Dr Géraldine Lavigne

Investigateur coordonnateur :
Dr Guy Aya

PHRC multicentrique

Promoteur : CHU Nîmes
(Nîmes, Clamart, Cochin, Lille, Nice, Montpellier, Genève, Saint Etienne)

42 patientes par groupe, soit 84 patientes

CJP : ⊙ 30% du taux d'embolisation ou de ligature vasculaire avec rFVIIa

ACJ : Nombre de sujets à traiter pour gagner une embolisation ou ligature, ...

Use of Recombinant Activated Factor VII in Primary Postpartum Hemorrhage

The Northern European Registry 2000–2004

Zarko Alfirevic, MD, FRCOG, Diana Elbourne, PhD, Sue Pavord, MBChB, FRCP, Antoinette Bolte, MD, Herman Van Geijn, MD, Frederic Mercier, MD, Jouni Ahonen, MD, Katarina Bremme, MD, Birgit Bødker, MD, Ebba Margrét Magnúsdóttir, MD, Kjell Salvesen, MD, Walter Prendiville, MAO, FRCOG, Ann Truesdale, BSc, Felicity Clemens, MPhil, Deborah Piercy, BSc, and Gill Gyte, BSc

1270 VOL. 110, NO. 6, DECEMBER 2007

OBSTETRICS & GYNECOLOGY

NEFOH

rVIIa $\approx$ 90 μ g/kg

108 patients (65 maternités / 6 pays)

Amélioration	83%
Identique ou pire	17%

- **Embolie Pulmonaire (2)**
- **Thrombose veineuse bilatérale 4 semaines plus tard**
- **IDM après arrêt cardiaque**
- **Eruption cutanée**

OFFHO : 2005-2007

Effets indésirables	Nombre de cas
Thrombose veineuse membre supérieur	1
Thrombose veineuse membre inférieur	3
Thrombose veineuse ovarienne	2
Embolie pulmonaire	3

103 CRF

rFVIIa (Novoseven[®]) et HPP :

Protocole Thérapeutique Temporaire (PTT)

AFSSAPS mars 2008 (pour 4 ans)

(<http://www.sante-ra.fr/doc/omedit/novoseven%20PTT%201.pdf>)

- après procédure invasive (ligatures vasc. ou embolisation) :
 - si l'HPP continue
 - avant hystérectomie si possible
- dose $\Rightarrow$ 90 $\mu\text{g}/\text{kg}$,
- renouvelable 1 fois (120 $\mu\text{g}/\text{kg}$)

+ normothermie, pH > 7,20, [Ca] $\sim$ Na^{al}
plaquettes > 30-50.000, fibrinogène > 0,5-1 g/L

Éditorial

Le rFVIIa dans le traitement de l'hémorragie du post-partum : évaluer le rapport bénéfique/risque

The role of rFVIIa in treatment of severe postpartum haemorrhage: To evaluate the risk/benefit ratio

Annales Françaises d'Anesthésie et de Réanimation 29 (2010) 673–675

A.-G. Aya

*Service d'anesthésie-réanimation, groupe hospitalier universitaire
Caremeau, université de Montpellier, 30029 Nîmes, France*

F.-J. Mercier*

*Département d'anesthésie-réanimation, hôpital A. Béclère,
université Paris-Sud, 157, rue de la porte-de-Trivaux,
BP 405, 92141 Clamart cedex, France*

*Auteur correspondant.

Adresses e-mail : frederic.mercier@abc.aphp.fr (F.-J. Mercier)

Use of clotting factors and other prohemostatic drugs for obstetric hemorrhage

Frédéric J. Mercier^a and Marie-Pierre Bonnet^{b,c}

^aDépartement d'Anesthésie Réanimation, Hôpital Antoine Bécclère, Groupe Hospitalier Universitaire Paris Sud, Assistance Publique-Hôpitaux de Paris, Université Paris XI, Clamart, ^bDépartement d'Anesthésie Réanimation, Maternité Port Royal, Hôpital Cochin, Groupement Hospitalier Universitaire Ouest, Assistance Publique-Hôpitaux de Paris, Université Paris V – René Descartes and ^cINSERM, UMR S953, Epidemiological Research Unit on Perinatal Health and Women's and Children's Health, Université Paris VI, Paris, France

Correspondence to Frédéric J. Mercier, Département d'Anesthésie-Réanimation Chirurgicale, Hôpital Antoine Bécclère, 157, rue de la Porte de Trivaux, 92141 Clamart Cedex BP 405, France
Tel: +33 1 45 37 42 73; fax: +33 1 45 37 49 85;
e-mail: frederic.mercier@abc.aphp.fr

Current Opinion in Anaesthesiology 2010, 23:000–000

Purpose of review

To guide the optimal use of blood products and to clarify the indications of prohemostatic drugs in obstetric hemorrhage.

Recent findings

The literature underlines the usefulness of transfusing packed red blood cells, fresh frozen plasma and platelets earlier and in defined ratios to prevent dilutional coagulopathy during obstetric hemorrhage. The value of fibrinogen concentrate and prohemostatic drugs such as tranexamic acid and recombinant factor VIIa is also pointed out.

Summary

It seems reasonable to use blood products transfusion earlier and in a 1 : 1 fresh frozen plasma : red blood cell ratio during acute obstetric hemorrhage; however, this analysis is mainly based on trauma literature. Fibrinogen concentrate should be added if fibrinogen plasma level remains below 1.0 g l^{-1} and maybe even as soon as it falls below $1.5\text{--}2.0 \text{ g l}^{-1}$; the addition of tranexamic acid (1 g) is cheap, likely to be useful and appears safe. Data on proactive administration of platelets are insufficient to recommend this practice routinely. Presently, recombinant factor VIIa ($60\text{--}90 \mu\text{g kg}^{-1}$) is advocated only after failure of other conventional therapies, including embolization or conservative surgery, but prior to obstetric hysterectomy. Prospective randomized controlled trials are highly desirable.

Pl. accreta

adhésion et/ou pénétration
anormale du Pl.

Arterial balloon occlusion of the internal iliac arteries for treatment of life-threatening massive postpartum haemorrhage: a series of 15 consecutive cases

J.P.M. Penninx^{a,*}, H.L.M. Pasmans^b, S.G. Oei^{a,c}

^a Department of Obstetrics and Gynecology, Máxima Medical Centre, Veldhoven, The Netherlands

^b Department of Radiology, Máxima Medical Centre, Veldhoven, The Netherlands

^c Department of Electrical Engineering, University of Technology, Eindhoven, The Netherlands

(BJOGRM 2009)

Fig. 1. Balloon occlusion of the internal iliac arteries.

Arterial balloon occlusion of the internal iliac arteries for treatment of life-threatening massive postpartum haemorrhage: a series of 15 consecutive cases

J.P.M. Penninx^{a,*}, H.L.M. Pasmans^b, S.G. Oei^{a,c}

^a Department of Obstetrics and Gynecology, Máxima Medical Centre, Veldhoven, The Netherlands

^b Department of Radiology, Máxima Medical Centre, Veldhoven, The Netherlands

^c Department of Electrical Engineering, University of Technology, Eindhoven, The Netherlands

(BJOGRM 2009)

Patients who underwent balloon occlusion because of massive life-threatening haemorrhage.

	Year	Gravity/ parity	Vaginal birth (V) or cesarean section (C)	Diagnosis	Blood loss (ml)	Packed cells	FFP ^a	Thrombocytes	Hysterectomy ^b	Place of procedure ^c
1	2007	G1P0	V	Retained placenta, uterine atony	5,000	10	2	1	No	OR
2	2007	G2P1	C	Uterine atony	7,000	10	4	0	No	OR
3	2007	G2P0	V	Uterine atony	8,000	14	8	1	No	OR
4	2006	G1P0	C	Uterine atony	7,000	15	3	0	No	OR
5	2006	G6P2	V	Retained placenta, uterine atony	5,000	10	3	0	No	OR
6	2006	G1P0	C	Uterine atony	10,000	16	4	1	No	OR
7	2005	G2P0	V	Uterine atony	6,000	16	6	2	No	OR
8	2005	G5P2	C	Placenta percreta, uterine atony	10,000	28	12	3	After	OR
9	2004	G1P0	C	Uterine atony	11,000	27	8	3	No	OR
10	2004	G2P0	V	Retained placenta, uterine atony, placenta accreta	10,000	22	8	2	After	OR
11	2004	G1P0	V	Uterine atony	5,600	7	4	0	No	OR
12	2003	G2P1	C	Placenta praevia, uterine atony, placenta increta	15,000	44	11	4	After	OR
13	1999	G3P2	V	Retained placenta, uterine atony	5,000	7	1	0	No	OR
14	1999	G2P1	V	Uterine atony, uterine rupture	15,000	69	33	8	Before	IC
15	1999	G2P1	C	Uterine atony	12,000	30	9	2	No	R

^a Fresh frozen plasma.

^b Hysterectomy performed before or after balloon occlusion.

^c IC, intensive care unit; R, radiology department; O, operating room.

Conservative management of PPH

- Preventive catheterization of the descending aorta via transhumeral access
- Stark cesarean delivery
- Uterotonic drugs
- Affronti endouterine square hemostatic sutures
- Preparation of B-Lynch suture
- Intrauterine application of Bakri balloon and partial filling with 100 mL of normal saline
- Perform the B-Lynch suture, hysterorrhaphy, and fill a Bakri balloon with up to 500 mL of normal saline
- Perform reversible radiological embolization and/or surgical ligation of the uterine arteries

Radical treatment of PPH

- Hysterectomy

Fig. 1. Protocol for the management of postpartum hemorrhage.

Table 1. Medical Management of Postpartum Hemorrhage

Table 2. Tamponade Techniques for Postpartum Hemorrhage

Technique	Comment
Uterine tamponade	
—Packing	—4-inch gauze; can soak with 5,000 units of thrombin in 5 mL of sterile saline
—Foley catheter	—Insert one or more bulbs; instill 60–80 mL of saline
—Sengstaken–Blakemore tube	
—SOS Bakri tamponade balloon	300–500 mL

Étude Pithagore 6.

	Oui	Non
Délivrance dirigée réalisée	61,7 %	38,3 %
Appel senior obstétricien-anesthésiste	68 %	32 %
Réalisation révision utérine	79 %	21 %
Injection supplémentaire ocytocine	93 %	7 %
Sulprostone administrée si indication	50 %	50 %
Sulprostone administrée < 30'	23 %	77 %
Bilan biologique < 60'	37 %	63 %
Hémorragie repérée en salle d'accouchement	68 %	32 %
Hémorragies du post-partum – Évaluation des pratiques professionnelles de prévention et de prise en charge – À propos de 106 000 accouchements et 9360 HPP		

19 janvier 2010 / n° 2-3

Numéro thématique - La mortalité maternelle en France : bilan 2001-2006
Special issue - Maternal deaths in France: situation report, 2001-2006

- ◆ Feuille de surveillance de l'HPP : modèle
- ◆ Après échec du Nalador, effectuer rapidement les mesures invasives des RPC
- ◆ Toutes les maternités doivent prévoir dans leur organisation la venue sur place d'un chirurgien (GO ou non), lorsque le MDG n'en a pas la compétence
- ◆ Toute structure doit être organisée pour effectuer les premiers gestes d'hémostase sur place quand la gravité de l'hémorragie l'impose

Resuscitation, monitoring, investigation and treatment should occur simultaneously

PREVENTION AND MANAGEMENT OF POSTPARTUM HAEMORRHAGE

This is the first edition of this guideline.

Guidelines for the use of rFVIIa in massive obstetric Haemorrhage

Welsh A et al, Aust N Z J Obstet Gynaecol 2008; 48: 12-16

Flow chart for management of PPH

INTERVENTIONS

(Notify local transfusion specialist of possible need for activation of Massive Blood Transfusion Protocol)

Medical:

- Treat: haemodynamic instability; hypothermia; acidosis
- Uterine massage / compression
- Uterotonic agents
- Coagulation studies and treat coagulopathy

Blood Component Therapy

- 4 U packed red blood cells (PRBCs)
 - Coagulopathy correction
 - 4 U PRBCs
 - 4 U FFP
 - Single adult dose of platelets
 - Repeat PRBCs, FFP and platelets
 - Administer calcium as appropriate
- Repeat (b) and (c) as necessary

Surgical (as available and appropriate)

- EUA and repair
- Uterine tamponade
- B-Lynch suture
- Arterial ligation
- Radiological arterial embolisation

Checklist for 'off-label' use of rFVIIa in obstetrics

- Remember the high risk of thromboembolism
- Consider physical measures for thromboprophylaxis
- Monitor all women for signs of improvement and adverse events.
- Report all patients receiving rFVIIa to the Haemostasis Registry (Monash University)

<http://www.med.monash.edu.au/epidemiology/traumaepi/haemostasis.html>

Figure 1 Flow chart for management of post-partum haemorrhage.

Major Obstetric Haemorrhage Flow Chart

MAJOR OBSTETRIC HAEMORRHAGE

TASK

Hémorragie en salle de naissance

Conclusion

- ◆ Les règles de base des RPC sur l'HPP de déc. 2004 demeurent valables
- ◆ Nouveautés de ces dernières années :
 - transfusion massive et les hémostatiques (ATX, fibri & rFVIIa)
 - prise en charge réanimatoire plus précoce et plus « agressive »
 - réflexions encore « embryonnaires » sur :
 - la place des ballons intra-utérins, notamment avant transfert, ou transport au bloc opératoire ou en embolisation
 - une stratégie spécifique pour l'HPP survenant durant la césarienne
 - une stratégie spécifique pour le P. accreta / percreta
 - la place des ballons en iliaques internes, voire en iliaques communes ou dans l'aorte sous-rénale

BEGAR 2011

Hôtel Pullman Montparnasse - PARIS

Mer 7 et Jeu 8 décembre :
→ Anesthésie-Réanimation Obstétricale

secret.anesthesieb@abc.aphp.fr

(ou 01 45 37 42 73)